

Exact Static Solutions for Scalar Fields Coupled to Gravity in $(3 + 1)$ -Dimensions

Ayşe H. Bilge and Durmus Daghan

Istanbul Technical University, Faculty of Science and Letters, Mathematics
Engineering Department, Maslak, TR-34469 Istanbul, TURKEY

E-mail: bilge@itu.edu.tr, daghand@itu.edu.tr

Abstract. Einstein's field equations for a spherically symmetric metric coupled to a massless scalar field are reduced to a system effectively of second order in time, in terms of the variables $\mu = m/r$ and $y = (\alpha/ra)$, where a , α , r and m are as in [W.M. Choptuik, "Universality and Scaling in Gravitational Collapse of Massless Scalar Field", *Physical Review Letters* **70** (1993), 9-12]. Solutions for which μ and y are time independent may arise either from scalar fields with $\phi_t = 0$ or with $\phi_s = 0$ but ϕ linear in t , called respectively the positive and negative branches having the Schwarzschild solution characterized by $\phi = 0$ and $\mu_s + \mu = 0$ in common. For the positive branch we obtain an exact solution which have been in fact obtained first in [I.Z. Fisher, "Scalar mesostatic field with regard for gravitational effects", *Zh. Eksp. Teor. Fiz.* **18** (1948), 636-640, gr-qc/9911008] and rediscovered many times (see D. Grumiller, "Quantum dilaton gravity in two dimensions with matter", PhD thesis, *Technische Universität, Wien* (2001), gr-qc/0105078) and we prove that the trivial solution $\mu = 0$ is a global attractor for the region $\mu_s + \mu > 0$, $\mu < 1/2$. For the negative branch discussed first in [M. Wyman, "Static spherically symmetric scalar fields in general relativity", *Physical Review D* **24** (1981), 839-841] perturbatively, we prove that $\mu = 0$ is a saddle point for the linearized system, but the non-vacuum solution $\mu = 1/4$ is a stable focus and a global attractor for the region $\mu_s + \mu > 0$, $\mu < 1/2$.

Keywords: Choptuik spacetime; Spherically symmetric static solutions; Massless scalar fields

PACS numbers: 83C15, 83C20

1. Introduction

The initial value problem for Einstein's equations with massless scalar field was studied analytically by Christodoulou in [1] and [2] where it was shown that "small initial data" disperses while for "large initial data" the end state is a black hole surrounded by vacuum. The work of Choptuik [3] on the numerical search for "critical initial data" that would separate these two types of behavior led to the discovery of the "threshold phenomenon". For various one parameter families of initial data, where p denotes the amplitude or the width of pulses of various shapes, it was observed that for $p \ll p_*$ the end state of the evolution is a flat space-time (external field dissipates), while for $p \gg p_*$ the end state of the evolution is a black hole. For p near p_* , where the black hole formation first starts, the black hole mass is proportional to $(p - p_*)^\gamma$ where the constant γ is nearly equal to 0.37 independently of the initial data. Furthermore, the critical solution exhibits a universal scaling behavior in the sense that it is periodical in

suitable time and space coordinates with a period nearly equal to 3.44 for all families. Threshold behavior was studied afterwards for a number of configurations and similar types of critical behavior have been observed. A detailed overview of the literature on the threshold phenomena can be found for example in [4].

The field equations used in numerical studies in the literature form a consistent but overdetermined system. In this paper we shall make use of the complete set of field equations as in [5] to decouple the equations for the metric functions from the equations for the scalar field. These equations form a normal system and they are effectively of second order in time. In logarithmic coordinates, they have no explicit dependency on the independent variables, as expected from the symmetry analysis of the original equations. The second order hyperbolic equation for the mass density involves a parameter that can be either plus or minus one, hence the decoupled system has positive and negative “branches”. Besides its suitability for perturbative studies, in the time independent case, this system reduces to an autonomous second order system and allows a rigorous analysis of the negative branch, which is not transparent in the original formulation.

When the field equations are written in that form, a non-flat, special solution for the negative branch is obvious: $\mu = \frac{1}{4}$, or $m = \frac{1}{4}r$, for which the external field is linear in time. Thus surprisingly, time varying external fields give rise to static metrics. A perturbative study of the full system is deferred to a future work. In Sections 4 and 5 we restrict ourselves to the static case, which can be realized either as the ones corresponding to time invariant external fields or the ones with $\phi_s = 0$ but $\phi_t \neq 0$. For the positive branch we obtain an exact analytic solution while for the negative branch, we prove that all solutions starting in the region $\mu_s + \mu > 0$, $\mu < 1/2$, stay in that region and the constant solution $\mu = \frac{1}{4}$ is a global attractor. Although there is no non-vacuum static black hole solution, for large values of ϕ_s or ϕ_t , μ gets very close to $1/2$ and decays rapidly to negative values, as depicted in Figure 3b. The trivial solution $\mu = 0$ and the Schwarzschild solution with $\mu_s + \mu = 0$ are of course common to the both branches.

It was pointed out by D. Grumiller in a private communication that the solution for the positive branch dates back to Fisher [6], as discussed in detail in [7] and [8], Fisher’s solution have been rediscovered in [9]-[13] and errors of the original equations 28-29 in [6] were corrected by Grumiller [7]. In addition we may cite the works of Bronnikov and Shikin [14] on the characterizations of the global properties of static, spherically symmetric configurations in various dimensions. As opposed to the popularity of the positive branch, the second class of static solutions that we call the negative branch is first noticed by Wyman [13] where it is studied perturbatively.

2. The Field Equations

Let M be a four dimensional Lorentzian manifold and g be a spherically symmetric metric on M . In the coordinate system adopted in [3], the local coordinates are $x^\mu = (t, r, \theta, \varphi)$ with

$$0 < \theta < \pi, \quad 0 < \varphi < 2\pi, \quad r > r_o > 0, \quad t > t_o > 0.$$

The metric is given by

$$ds^2 = -\alpha^2 dt^2 + a^2 dr^2 + r^2 d\theta^2 + r^2 \sin^2\theta d\varphi^2 \quad (2.1)$$

where $\alpha = \alpha(r, t)$, $a = a(r, t)$.

The Einstein's equations coupled to a scalar, static particle are [15]

$$R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R = \kappa T_{\mu\nu},$$

where

$$T_{\mu\nu} = \partial_\mu \phi \partial_\nu \phi - \frac{1}{2}g_{\mu\nu}(\partial_\lambda \phi \partial^\lambda \phi)$$

and ϕ satisfies the wave equation

$$g^{\mu\nu} \nabla_\mu \nabla_\nu \phi = (-g)^{-\frac{1}{2}} \partial_\mu [(-g)^{\frac{1}{2}} g^{\mu\nu} \partial_\nu \phi]$$

with $g = \det(g_{\mu\nu})$ [16]. The complete set of field equations are given below, where we have written $\kappa = 8\pi$ and omitted an equation which was identically satisfied.

$$\frac{a_t}{a} = 4\pi r \phi_t \phi_r, \quad (2.2a)$$

$$\frac{a_r}{a} - \frac{\alpha_r}{\alpha} + \frac{a^2 - 1}{r} = 0, \quad (2.2b)$$

$$\frac{a_r}{a} + \frac{\alpha_r}{\alpha} = 4\pi r \left[\phi_r^2 + \frac{a^2}{\alpha^2} \phi_t^2 \right], \quad (2.2c)$$

$$\left(\frac{a}{\alpha} \phi_t \right)_t = \frac{1}{r^2} \left(r^2 \frac{\alpha}{a} \phi_r \right)_r. \quad (2.2d)$$

Remark 2.1. The system above is over-determined but integrable in the sense that the equations for a_t and a_r are compatible, as noted in [5]. The degree of freedom in the system appears to be three functions of a single variable, corresponding to the initial data for ϕ , ϕ_t and α . This last freedom is actually redundant, since it can be eliminated by redefining t . The use of the complete system will allow the decoupling of the equations for a and α from the equation for ϕ .

The literature on the threshold phenomenon works with the system of equations (2.2b-d), starting with initial data for ϕ . We shall use the function $m(r, t)$ defined in terms of $a(r, t)$ as our primary variable and introduce logarithmic coordinates to obtain a normal (i.e. not overdetermined) system with no explicit dependency on the independent variables.

We start by defining the variables

$$z = \frac{\alpha}{a}, \quad a^2 = \left[1 - \frac{2m}{r} \right]^{-1}. \quad (2.3)$$

Note that as $r > 0$, we need to impose $r - 2m > 0$ to ensure the positivity of a^2 . The equations (2.2a-d) can be rearranged in the form

$$\frac{z_r}{z} = \frac{2m}{r(r - 2m)}, \quad (2.4a)$$

$$\frac{m_t}{r(r - 2m)} = 4\pi \phi_t \phi_r, \quad (2.4b)$$

$$\frac{m_r}{r(r-2m)} = 2\pi \left[\phi_r^2 + \frac{\phi_t^2}{z^2} \right], \quad (2.4c)$$

$$\left(\frac{1}{z} \phi_t \right)_t = \frac{1}{r^2} (r^2 z \phi_r)_r. \quad (2.4d)$$

Finally passing to logarithmic coordinates we obtain an autonomous system, as given below in Proposition 2.1.

Proposition 2.1. *Let*

$$s = \ln(r), \quad z = ry, \quad m = r\mu. \quad (2.5)$$

Then, the system of equations (2.2a-d) is equivalent to the system of equations

$$\frac{y_s}{y} = -2 + \frac{1}{1-2\mu}, \quad (2.6a)$$

$$\frac{1}{2\pi} \frac{\mu_t}{1-2\mu} = 2\phi_s \phi_t, \quad (2.6b)$$

$$\frac{1}{2\pi} \frac{\mu_s + \mu}{1-2\mu} = \phi_s^2 + \frac{\phi_t^2}{y^2}, \quad (2.6c)$$

$$\left(\frac{\phi_t}{y} \right)_t = 2y\phi_s + (y\phi_s)_s, \quad (2.6d)$$

where $\mu_s + \mu$ and μ_t/y are constrained by

$$\frac{\mu_s + \mu}{1-2\mu} \geq 0, \quad \frac{1}{2\pi} \frac{(\mu_s + \mu) \pm (\mu_t/y)}{1-2\mu} = \left(\phi_s \pm \frac{\phi_t}{y} \right)^2 \geq 0. \quad (2.7)$$

Remark 2.2. In addition to the translation invariance, the system (2.6a-d) admits the symmetry

$$(t, y) \rightarrow \left(\tau = \tau(t), \tilde{y} = \frac{1}{d\tau/dt} y \right). \quad (2.8)$$

Thus y can be scaled by an arbitrary function of t only, hence its time evolution is determined in terms of the time evolution of μ .

Remark 2.3. A special case of (2.8) is the transformation

$$(t, y) \rightarrow (\tau = \ln t, \tilde{y} = ty). \quad (2.9)$$

Hence one can use $(\tau, \tilde{y})$ as well as (t, y) in (2.6a-d). If $\tilde{y}$ is periodic in τ , i.e., $\tilde{y}(\tau, s) = \tilde{y}(\tau + T, s)$. Then

$$e^\tau y(e^\tau, s) = e^T e^\tau y(e^T e^\tau, s),$$

or

$$y(t, s) = e^T y(e^T t, s),$$

hence periodic solutions lead to the “echoes” of the original solution.

The regions determined by the conditions $\phi_s = \pm \frac{\phi_t}{y}$ and $\mu_s + \mu = \pm \frac{\mu_t}{y}$ are shown in the Figures 1a and 1b respectively. As the equations (2.6a-d) are invariant under

the discrete symmetry $\phi_s \rightarrow -\phi_s$ it is sufficient to work with ϕ in regions *I* and *II* for example. Also for $1 - 2\mu > 0$ only region I^* in Figure 1b is allowed. As it will be seen in the next section, the half plane consisting of the regions *I* and *II* in the $(\phi_s, \phi_t/y)$ plane is mapped to the quarter plane I^* as a double cover. We shall call these as the “positive” and “negative” branches. The boundary of the regions *I* and *II* is characterized by $\phi_s = \pm \phi_t/y$ or equivalently with $\mu_s + \mu = \pm \mu_t/y$. The decoupled equation (3.1b) below admits non-vacuum solutions $\mu_s + \mu = \pm \mu_t/y$, but when we take into account the wave equation for ϕ we get $\phi_s = \phi_t/y = 0$, hence $\mu_s + \mu = \mu_t = 0$.

3. Reduction of the field equations

We shall eliminate ϕ from (2.6a-d) and obtain a system of equations for μ and y . The proof of the proposition below is essentially a straightforward computation. We have introduced the auxiliary variables X and Y below in order to make the proof more readable.

Proposition 3.1. *For $\phi_s^2 - (\frac{\phi_t}{y})^2 \neq 0$ the system of equations (2.6a-d) is equivalent to the system of equations given below.*

$$\frac{y_s}{y} = -2 + \frac{1}{(1 - 2\mu)}, \quad (3.1a)$$

$$\begin{aligned} \mu_{ss} + \mu_s - \frac{1}{y} \left(\frac{\mu_t}{y} \right)_t + \frac{2}{(1 - 2\mu)} \left[(\mu_s + \mu)^2 + (\mu_s + \mu)\mu - \frac{\mu_t^2}{y^2} \right] \\ \pm 2\epsilon \left[(\mu_s + \mu)^2 - \frac{\mu_t^2}{y^2} \right]^{1/2} = 0, \end{aligned} \quad (3.1b)$$

$$\phi_s^2 + (\phi_t/y)^2 = \frac{1}{2\pi} \frac{\mu_s + \mu}{1 - 2\mu}, \quad (3.1c)$$

$$2\phi_s(\phi_t/y) = \frac{1}{2\pi} \frac{\mu_t/y}{1 - 2\mu}. \quad (3.1d)$$

where $\epsilon = \text{sgn}(1 - 2\mu)$.

Proof. Let

$$X = \phi_s^2 + (\phi_t/y)^2 = \frac{1}{2\pi} \frac{\mu_s + \mu}{1 - 2\mu},$$

$$Y = 2\phi_s(\phi_t/y) = \frac{1}{2\pi} \frac{\mu_t/y}{1 - 2\mu}.$$

It can be seen that either of the conditions $\phi_s = \pm \frac{\phi_t}{y}$ is equivalent to $\sqrt{X^2 - Y^2} = 0$. In the following we assume that $X^2 - Y^2 \neq 0$. Note that $\mu_t = 0$ hence $Y = 0$ can be achieved either by taking $\phi_s = 0$ or $\phi_t = 0$. These time independent solutions will be discussed in Section 4. Assuming for example $\phi_s \neq 0$, we obtain

$$\phi_t/y = \frac{Y}{2\phi_s}.$$

Substituting this in X we have

$$\phi_s^4 - X\phi_s^2 + \frac{Y^2}{4} = 0.$$

From this we can solve

$$\phi_s^2 = \frac{1}{2}[X \pm \sqrt{X^2 - Y^2}].$$

Writing

$$\Delta = X \pm \sqrt{X^2 - Y^2}$$

we have

$$\phi_s = \pm \frac{1}{\sqrt{2}}\Delta^{1/2}, \quad \phi_t/y = \pm \frac{1}{\sqrt{2}}Y\Delta^{-1/2}.$$

As noted at the end of Section 2, the signatures of ϕ_s and ϕ_t are irrelevant, hence we may take the positive signature.

After substituting these and multiplying with $\Delta^{1/2}$, the differential equation for ϕ reduces to

$$\frac{1}{y} \left[Y_t - \frac{1}{2}Y \frac{\Delta_t}{\Delta} \right] = \frac{1}{1-2\mu}\Delta + \frac{1}{2}\Delta_s.$$

Substituting Δ and taking derivatives one can see that the coefficients of X_s and Y_t are proportional, in fact we have

$$\frac{1}{2}[1 \pm X(X^2 - Y^2)^{-1/2}][Y_t/y - X_s] = \frac{1}{1-2\mu}(X \pm \sqrt{X^2 - Y^2}) \mp \frac{1}{2} \frac{Y(Y_s - X_t/y)}{(X^2 - Y^2)^{1/2}}.$$

It can be seen that

$$Y_s - X_t/y = -\frac{4\mu}{1-2\mu}Y.$$

Hence it follows that

$$X_s - Y_t/y + \frac{4\mu}{1-2\mu}X \pm 2\sqrt{X^2 - Y^2} = 0$$

which is (3.1b). •

For the rest of the paper we assume that $\mu_s + \mu > 0$, hence we take $\epsilon = 1$ in the equation (3.1b).

4. Static metrics: Exact solution for the positive branch

In this section we shall study solutions for which the metric is independent of t , hence static. This will allow nevertheless a time dependency in the scalar field. Putting $\mu_t = 0$, the system of equations (3.1a-d) reduces to

$$\frac{y_s}{y} = -2 + \frac{1}{1-2\mu}, \tag{4.1a}$$

$$\mu_{ss} + \mu_s + \frac{2}{1-2\mu}(\mu_s + \mu)(\mu_s + 2\mu) \pm 2(\mu_s + \mu) = 0, \tag{4.1b}$$

$$\phi_s^2 + (\phi_t/y)^2 = \frac{1}{2\pi} \frac{\mu_s + \mu}{1-2\mu}, \tag{4.1c}$$

$$2\phi_s(\phi_t/y) = \frac{1}{2\pi} \frac{\mu_t/y}{1-2\mu}. \tag{4.1d}$$

We shall first give special solutions with $\mu_s = k\mu$, then obtain the exact solution for the positive branch and finally study the solutions for the negative branch in Section 5.

Special solutions with $\mu_s = k\mu$.

For the positive and negative branches we obtain respectively,

$$\text{positive branch: } (k+1)(k+2)\mu = 0,$$

$$\text{negative branch: } (k+1)[8\mu^2 + (k-2)\mu] = 0.$$

Exact solutions for these special cases are given below with equations (4.2-5), where y_o and μ_o are integration constants.

1. $\mu = 0$ is a special solution for both positive and negative branches for which $\phi_s = \phi_t = 0$ and $y = y_o e^{-s}$. The line element is the flat metric

$$ds^2 = -y_o^2 dt^2 + dr^2 + r^2 d\theta^2 + r^2 \sin^2\theta d\varphi^2. \quad (4.2)$$

2. $\mu = 1/4$ is a solution for the negative branch corresponding to $\phi_s = 0$ and $\phi_t/y = 1/2\sqrt{\pi}$. For this case y is a constant. Note that such a solution intersects the $m = r/2$ line only at the origin. Thus its perturbations may lead to what is observed as black holes with infinitesimal mass. The metric (2.1) becomes

$$ds^2 = -2y_o^2 r^2 dt^2 + 2 dr^2 + r^2 d\theta^2 + r^2 \sin^2\theta d\varphi^2. \quad (4.3)$$

The only nonzero component of the Ricci tensor is $R_{tt} = 2y_o^2$ and the Ricci scalar is $R = -\frac{1}{r^2}$.

3. $k = -1$, or $\mu_s + \mu = 0$ corresponding to constant m is a solution for positive and negative branches. Here $\mu = \mu_o e^{-s}$ and $y = \frac{y_o}{\mu_o} \mu |1 - 2\mu|$. This is the Schwarzschild solution, it is the only possibility for intersecting the $\mu = 1/2$ line. For increasing s the solution curves tend to the origin. For negative μ they escape to infinity in the reverse direction. When μ_o is positive, it has the standard interpretation as the black hole mass. The metric (2.1) becomes

$$ds^2 = -y_o^2 \left[1 - \frac{2\mu_o}{r} \right] dt^2 + \left[1 - \frac{2\mu_o}{r} \right]^{-1} dr^2 + r^2 d\theta^2 + r^2 \sin^2\theta d\varphi^2. \quad (4.4)$$

4. $k = -2$ or $\mu_s + 2\mu = 0$ is a solution for the positive branch. Then $\mu = \mu_o e^{-2s}$ and $y = y_o \left| \frac{\mu}{\mu_o} \right|^{1/2} |1 - 2\mu|^{1/2}$. For increasing s the curves tend to the origin. The part of the curve for positive μ is not allowed. The part for negative μ escapes to infinity in the reverse direction. The metric (2.1) becomes

$$ds^2 = -y_o^2 dt^2 + \left[1 - \frac{2\mu_o}{r^2} \right]^{-1} dr^2 + r^2 d\theta^2 + r^2 \sin^2\theta d\varphi^2. \quad (4.5)$$

The only nonzero component of the Ricci tensor is $R_{rr} = \frac{4\mu_o}{r^2(2\mu_o - r^2)}$ and Ricci scalar is $R = -\frac{4\mu_o}{r^4}$.

Exact solution for the positive branch.

For the positive branch $\phi_t = 0$ and we write $\psi = \phi_s$. The field equations reduce to

$$\frac{y_s}{y} = -2 + \frac{1}{1 - 2\mu} = \frac{4\mu - 1}{1 - 2\mu}, \quad (4.6a)$$

$$\mu_s = -\mu + 2\pi(1 - 2\mu)\psi^2, \quad (4.6b)$$

$$\frac{\psi_s}{\psi} = -\frac{1}{1 - 2\mu}. \quad (4.6c)$$

We will obtain an analytic solution for this system.

Proposition 4.1. For $\phi_t = 0$ the solution of the system (4.6a-c) is

$$\mu = \frac{1}{2} \left[\frac{\psi^2 - \frac{c}{4\pi}\psi}{\psi^2 - \frac{c}{4\pi}\psi - \frac{1}{4\pi}} \right], \quad (4.7)$$

$$\frac{y}{y_o} = |\psi| |\psi - p|^{B_1} |\psi + q|^{B_2}, \quad (4.8)$$

$$\frac{r}{r_o} = |\psi|^{-1} |\psi - p|^{C_1} |\psi + q|^{C_2} \quad (4.9)$$

where $\psi = \phi_s$, $r = e^s$, y_o and r_o are constants,

$$p = \frac{1}{8\pi} \left(c + \sqrt{c^2 + 16\pi} \right) = \frac{c}{8\pi} \left(1 + \frac{1}{\delta} \right), \quad (4.10a)$$

$$q = \frac{1}{8\pi} \left(-c + \sqrt{c^2 + 16\pi} \right) = \frac{c}{8\pi} \left(-1 + \frac{1}{\delta} \right), \quad (4.10b)$$

$$B_1 = -(1 - \delta), \quad B_2 = -(1 + \delta), \quad C_1 = \frac{1}{2}(1 - \delta), \quad C_2 = \frac{1}{2}(1 + \delta), \quad (4.10c)$$

where

$$\delta = \frac{c}{\sqrt{c^2 + 16\pi}}. \quad (4.11)$$

Proof. Let $u = (1 - 2\mu)\psi$. Taking the derivative of u with respect to ψ and using (4.6b,c), we get

$$\frac{du}{d\psi} = u^2 \left(4\pi + \frac{1}{\psi^2} \right).$$

This is a separable equation which can be solved as

$$u = \frac{\psi}{1 + c\psi - 4\pi\psi^2}$$

where c is an integration constant. From the definition of u we obtain μ as a function of ψ as in (4.7). Note that there are vertical asymptotes at $\psi = p$ and $\psi = -q$, with p and q as in (4.10a,b). In the region $-q < \psi < p$, μ is less than $1/2$ and has a unique maximum at $\psi_M = c/8\pi$ where

$$\mu_M = \mu(\psi_M) = \mu(c/8\pi) = \frac{1}{2} \frac{c^2}{c^2 + 16\pi} = \frac{1}{2} \delta^2. \quad (4.12)$$

For $|c| > 4\sqrt{\pi}$, μ crosses the level $\mu = 1/2$ twice at $\psi^\pm$ given by

$$\psi^\pm = \frac{1}{8\pi} \left(c \pm \sqrt{c^2 - 16\pi} \right). \quad (4.13)$$

A typical solution curve is given in Figure 2. Note that

$$\frac{d\mu}{d\psi} \big|_{\psi=0} = \frac{1}{2}c,$$

i.e., the parameter c is the slope of the curve $\mu(\psi)$ at $\psi = 0$ or as $s \rightarrow \infty$. As we may assume $\psi > 0$, we shall restrict ψ in the range $0 < \psi < p$. For c negative and zero, μ is negative, ranging from $-\infty$ to 0 as ψ changes from p to 0. For $0 < c < 4\sqrt{\pi}$ it has a maximum at ψ_M but lies below the $\mu = 1/4$ line. As c gets large, the solution curve is nearly a broken line following closely the borders of the region bounded by $\mu = 1/2$, $\psi = p$ and $\psi = -q$. For the critical value of $c = 4\sqrt{\pi}$, $\mu_M = 1/4$ at $\psi_M = 1/2\sqrt{\pi}$ and $p = \frac{1}{2\sqrt{\pi}}(-1 + \sqrt{2})$. Typical solution curves are presented in Figure 3a.

To solve for y we write

$$\frac{dy}{y} = \frac{d\psi}{\psi} (1 - 4\mu) = \frac{d\psi}{\psi} \frac{[-\psi^2 + \frac{c}{4\pi}\psi - \frac{1}{4\pi}]}{[\psi^2 - \frac{c}{4\pi}\psi - \frac{1}{4\pi}]} = d\psi \left[\frac{A}{\psi} + \frac{B_1}{\psi - p} + \frac{B_2}{\psi + q} \right]$$

where $A = 1$ and B_1 and B_2 are as in (4.10c). By integrating we obtain y as in (4.8). It can be seen that $dy/d\psi$ vanishes at $\psi^\pm$ for which $\mu(\psi^\pm) = 1/4$. Hence for $|c| \leq 4\sqrt{\pi}$, y is a monotone increasing function of ψ while for $|c| > 4\sqrt{\pi}$, y has two critical points. As it can be seen from Figure 4, for large c , the variation of y with ψ is quite sharp.

Finally, substituting the expression of μ in (4.6c), we obtain the equation

$$\frac{d\psi}{4\pi\psi(\psi^2 - \frac{c}{4\pi}\psi - \frac{1}{4\pi})} = d\psi \left[\frac{A}{\psi} + \frac{C_1}{\psi - p} + \frac{C_2}{\psi + q} \right] = ds$$

where $A = -1$ and C_1 and C_2 are given by (4.10c). Integrating this and exponentiating we obtain (4.9). It can be seen that $r = e^s$ is a monotone decreasing function of ψ . Parametric plots of μ , y and r in terms of ψ , are shown in Figure 4. •

From the remark above on the interpretation of c and a qualitative study of the solutions we see that the rate of change of the mass distribution with respect to ψ as $s \rightarrow \infty$ determines the solution completely. For large c , there is nearly a black hole formation at s^* determined by $\psi(s^*) = \psi_M = c/8\pi$, in the sense that, the solution curve nearly touches the $\mu = 1/2$ line but moves away from it to go eventually to large negative values as $s \rightarrow -\infty$ or as $r \rightarrow 0$.

We were unable to find an exact solution for the negative branch. For this case we shall study the equation (4.1b) as a dynamical system. To compare both branches and to analyze the efficiency of numerical algorithms, we also studied the positive branch of (4.1b) as a dynamical system in the μ - μ_s plane. Writing $\nu = \mu_s$, we can express the positive branch of (4.1b)

$$\begin{bmatrix} \mu_s \\ \nu_s \end{bmatrix} = \begin{bmatrix} \nu \\ -3\nu - 2\mu - \frac{2}{1-2\mu}(\nu^2 + 3\mu\nu + 2\mu^2) \end{bmatrix}. \quad (4.14)$$

The only critical point is $(0, 0)$ and the linearized part at the origin is

$$\begin{bmatrix} \bar{\mu} \\ \bar{\nu} \end{bmatrix}_s = \begin{bmatrix} 0 & 1 \\ -2 & -3 \end{bmatrix} \begin{bmatrix} \bar{\mu} \\ \bar{\nu} \end{bmatrix}. \quad (4.15)$$

The origin is a node and $\mu_s + \mu = 0$ and $\mu_s + 2\mu = 0$ are exact solutions. Typical solution curves shown in Figure 5 are sketched using the exact solution. Since ψ is a monotone function of s , one can as well think of the phase plane curves as parameterized by ψ and make parametric plots of μ and μ_s . It can be seen that curves in the second quadrant, lying between $\mu_s + \mu = 0$ and $\mu_s + 2\mu = 0$ correspond to c ranging from $-\infty$ to 0, $\mu_s + 2\mu = 0$ being associated with $c = 0$. The curves beyond $\mu_s + 2\mu = 0$ cross the $\mu = 0$ line at a finite value of the parameter and reach the origin as $s \rightarrow \infty$. The expressions of μ and μ_s obtained from the exact solution implies that the origin is a global attractor for all solutions starting in the region defined by $\mu_s + \mu > 0$ and $\mu < 1/2$.

5. Static metrics: Phase plane analysis for the negative branch

Writing $\nu = \mu_s$, we can express second order equation (4.1b) for μ as a dynamical system

$$\begin{bmatrix} \mu_s \\ \nu_s \end{bmatrix} = \begin{bmatrix} \nu \\ \nu + 2\mu - \frac{2}{1-2\mu}(\nu^2 + 3\mu\nu + 2\mu^2) \end{bmatrix}, \quad (5.1)$$

where $\nu = \mu_s$. There are two critical points, $(0, 0)$ and $(1/4, 0)$. The linearized part at the origin is

$$\begin{bmatrix} \bar{\mu} \\ \bar{\nu} \end{bmatrix}_s = \begin{bmatrix} 0 & 1 \\ 2 & 1 \end{bmatrix} \begin{bmatrix} \bar{\mu} \\ \bar{\nu} \end{bmatrix}. \quad (5.2)$$

To study the local behavior at $(0, 0)$ we note that the eigenvalues of the system above are -1 and 2 hence the origin is a saddle point. The corresponding eigenvectors are $(1, -1)$ and $(1, 2)$, hence the solutions curves approaching the origin along $\nu = -\mu$ in either direction are repelled and move away along $\nu = 2\mu$.

The linearized part at the point $(1/4, 0)$ is

$$\begin{bmatrix} \bar{\mu} \\ \bar{\nu} \end{bmatrix}_s = \begin{bmatrix} 0 & 1 \\ -4 & -2 \end{bmatrix} \begin{bmatrix} \bar{\mu} \\ \bar{\nu} \end{bmatrix}. \quad (5.3)$$

The eigenvalues of the system above are $-1 \pm i\sqrt{3}$, hence the point $(1/4, 0)$ is a stable focus. The local behavior near the points $(0, 0)$ and $(1/4, 0)$ will be crucial in the proof of the global behavior. We will now prove that $(1/4, 0)$ is a global attractor for solutions in the open half plane $\mu_s + \mu > 0$, $\mu < 1/2$.

Proposition 5.1. *Let (μ, ν) be a solution of (5.1) and D be the region bounded by $\mu + \nu > 0$ and $\mu < 1/2$. If $(\mu(0), \nu(0))$ belongs to D , then the solution curve $(\mu(s), \nu(s))$ remains in D for all s and*

$$\lim_{s \rightarrow \infty} (\mu(s), \nu(s)) = (1/4, 0).$$

Proof. Let V be the velocity vector of the solution curve. We can express V as

$$V = \left[\nu, -\frac{2}{(1-2\mu)}(\nu + 2\mu)(\nu + 2\mu - \frac{1}{2}) \right]. \quad (5.4)$$

As $1 - 2\mu$ is positive on D , the direction of V is determined by the signatures of ν , $\nu + 2\mu$ and $\nu + 2\mu - 1/2$. This leads to a subdivision of D into five regions as given below.

Region		μ_s	ν_s
D_1 :	$\nu > 0, \quad \mu + \nu > 0, \quad 2\mu + \nu < 0.$	> 0	< 0
D_2 :	$\nu > 0, \quad 2\mu + \nu \geq 0, \quad 1 - 4\mu - 2\nu < 0.$	> 0	> 0
D_3 :	$\nu > 0, \quad 1 - 4\mu - 2\nu \geq 0, \quad \mu < 1/2.$	> 0	< 0
D_4 :	$\nu \leq 0, \quad 1 - 4\mu - 2\nu > 0, \quad \mu < 1/2.$	≤ 0	< 0
D_5 :	$\nu \leq 0, \quad \mu + \nu > 0, \quad 1 - 4\mu - 2\nu \leq 0, \quad \mu < \frac{1}{2}.$	≤ 0	> 0

In the region D_1 , the velocity vector of the solution curve lies in the fourth quadrant. By uniqueness it cannot intersect the $\mu + \nu = 0$ line. Furthermore as the origin is a saddle point the solution curve necessarily intersects the $\nu + 2\mu = 0$ line and enters in the region D_2 .

The velocity vector of the solution curves in the region D_2 lie in the second quadrant, hence each solution curve passes to the region D_3 for increasing s .

In the region D_3 , the velocity vector is in the fourth quadrant. Here we will show that the boundaries $\nu + 2\mu = 1/2$ and $\mu = 1/2$ are never hit.

In order to show that solution curves do not hit the boundary $\nu = -2\mu + 1/2$, we prove that the slope of their velocity vector is larger than -2 . For this we need the inequality

$$(\nu + 2\mu)(\nu + 2\mu - 1/2) < \nu(1 - 2\mu)$$

to be satisfied. The equality in the relation above defines a hyperbola, the half line $\nu + 2\mu = 1/2$, $\mu < 1/4$ lies in the region where the inequality is satisfied. Hence away from the $\nu = 0$ level, the velocity vector V is surely pointing away from the $\nu + 2\mu = 1/2$ line. On the other hand as $(1/4, 0)$ is a focus, the solution curves cannot hit this point as ν gets close to zero, hence necessarily, cross the vertical line $\mu = 1/4$.

To prove that solution curves do not hit the line $\mu = 1/2$ at a point (μ, ν) with $\mu > 1/4$, we need to show that the velocity vector V points below the line joining (μ, ν) and $(1/2, 0)$, i.e., we need to prove that

$$|\nu_s/\mu_s| > |\nu/(\frac{1}{2} - \mu)|.$$

Since $(1 - 2\mu)$ and $\mu_s = \nu$ are positive in this region, we should have

$$|\nu_s| (1 - 2\mu) - 2\nu^2 > 0.$$

Substituting for ν_s , we obtain

$$4\mu\nu + (2\nu + 2\mu)(4\mu - 1) > 0.$$

For $\mu > 1/4$ this inequality is surely satisfied and solution curves reach the μ axis without hitting the $\mu = 1/2$ line.

In the region D_4 , the velocity vector belongs to the third quadrant, hence the solution curves hit the $\nu + 2\mu = 1/2$ line and pass to the region D_5 . In the region D_5 the velocity vector belongs to the second quadrant. By uniqueness of solutions, they cannot cross the $\mu + \nu = 0$ line and the local behavior guarantees that they stay away

from the origin and cross the μ axis. Again by uniqueness the solution curves spiral around the point $(1/4, 0)$ which is a stable focus. •

The integral curves for the numerical solution of the dynamical system 5.1 in the domain $\mu_s + \mu > 0$ and $\mu < 1/2$ are given in Figure 7. Expect for the line $\mu_s + \mu = 0$, all solution curves in this region spiral around the point $(1/4, 0)$ as $s \rightarrow \infty$. As one moves to right, the solution curves get very close to the line $\mu = 1/2$ but decay quickly to the level $\mu = 1/4$. The variations of μ , y and ϕ_t/y with respect to s exhibit rapidly decaying oscillations as shown in Figure 8.

The period of the linearized part of the dynamical system (5.1) at the point $(\frac{1}{4}, 0)$ is $\frac{2\pi}{\sqrt{3}} \simeq 3.62$. For the nonlinear system we could observe only 3 periods of oscillations. Starting with the initial conditions $(\mu = 0, \mu_s = \mu_s^0)$, μ_s^0 ranging from 10^{-7} to 10^3 , we computed consecutive crossings of the integral curves across $\mu = 1/4$ line at $\mu_s = a_i$, $i = 1, \dots, 4$ and we evaluated the periods T_i , $i = 1, 2, 3$ as the distance s between two consecutive crossings. Representative values for a_i 's and T_i 's are listed below.

$\mu_s^0 = 10^{-7} :$	a_1, a_2, a_3	0.0923	0.0022	0.0001
	T_1, T_2, T_3	3.5350	3.6280	3.6070
$\mu_s^0 = 1 :$	a_1, a_2, a_3	0.4698	0.0089	0.0002
	T_1, T_2, T_3	3.3510	3.6180	3.6380
$\mu_s^0 = 10^3 :$	a_1, a_2, a_3	567.4564	0.0837	0.0020
	T_1, T_2, T_3	4.7040	3.5430	3.6280

As μ_s^0 goes to infinity the first crossing level a_1 goes to infinity i.e., $\lim_{\mu_s^0 \rightarrow \infty} a_1 = \infty$, while for $i > 1$ they tend to a limit that we denote a_i^* , with

$$\lim_{\mu_s^0 \rightarrow \infty} a_{i+1} = a_i^* = \lim_{\mu_s^0 \rightarrow 0} a_i.$$

For $i = 1, 2, 3$, we have $a_1^* \simeq 0.092$, $a_2^* \simeq 0.0022$, $a_3^* \simeq 0.0001$.

Although the first oscillation depends on the initial condition the second oscillation start near $\mu_s = 0.0022 - 0.0090$ and have a quite stable period of 3.62, close to the period of the linearized system.

Remark 5.1. Putting $\mu_{tt} = k\mu_s$ in the linearized part of the full system and writing $\mu = \frac{1}{4} + u$ we have

$$u_{ss} + (2 - k)u_s + 4u = 0.$$

For $k = 2$, the solution of linearized and nonlinear system are unrelated as predicted by Hartman-Grobman theorem [17]. For arbitrary $0 < k < 2$ the linearized solution is of the form $u = e^{-\alpha s} e^{\pm i\omega s}$ where $\alpha = 1 - \frac{k}{2}$ and $\omega^2 + \alpha^2 = 4$. As an interesting coincidence we observed that the period corresponding to $k = 0.3704$ is $T = \frac{2\pi}{1.8265} = 3.44$.

Acknowledgments

The authors would like to thank Dr. D. Grumiller for the pointing out references [6]-[14] and Professor B.K. Harrison for illuminating discussions. This work is partially supported by the Turkish National Council for Scientific and Technological Research.

References

- [1] Christodoulou, D., The problem of a self-gravitating scalar field, *Commun. Math. Phys.* **105** (1986), 337-361.
- [2] Christodoulou, D., Global existence of generalized solutions of the spherically symmetric Einstein- scalar equations in the large, *Commun. Math. Phys.* **106** (1986), 587-621.
- [3] Choptuik, W.M., Universality and scaling in gravitational collapse of massless scalar field, *Physical Review Letters* **70** (1993), 9-12 .
- [4] Gundlach, C., Critical phenomena in gravitainal collapse, *Phys. Rept.* **376** (2003), 339-405, gr-qc/0210101.
- [5] Harrison, B.K., Applications of symmetry to general Relativity, *The fifty International Conference on Symmetry in Nonlinear Mathematical Physics*, **Part 1** (2004), 131-141.
- [6] Fisher, I.Z., Scalar mesostatic field with regard for gravitational effects, *Zh. Eksp. Teor. Fiz.* **18** (1948), 636-640, gr-qc/9911008.
- [7] Grumiller, D., Quantum dilaton gravity in two dimensions with matter, PhD thesis, *Technische Universität, Wien* (2001), gr-qc/0105078.
- [8] Grumiller, D., Mayerhofer, D., On static solutions in 2D dilaton gravity with scalar matter , *Class. and Quant. Grav.* **21** (2004), 5893-5914, gr-qc/0404013.
- [9] Bergmann, O., Leipnik, R., Space-time structure of a static spherically symmetric scalar field, *Phy. Rev.* **107** (1957), 1157.
- [10] Buchdahl, H.A, Reciprocal static metrics and scalar fields in the general theory of relativity, *Phy. Rev.* **115** (1959), 1325.
- [11] Janis, A.I., Newman, E.T., Winicour, J., Relativity of the Schwarzschild singularity, *Phys. Rev. Lett.* **20** (1968), 878.
- [12] Katanaev, M.O., Effective action for scalar fields in two-dimensional gravity, *Annals Phys.* **296** (2002), 1-50, gr-qc/0101033.
- [13] Wyman, M., Static spherically symmetric scalar fields in general relativity, *Physical Review D* **24** (1981), 839-841.
- [14] Bronnikov, K.A., Shikin, G.N., Spherically symmetric scalar vacuum: no-go theorems, black holes and solitons, *Grav. Cosmol.* **8** (2002) , 107-116, gr-qc/0109027.
- [15] Wald, R.M., General Relativity, *The University of Chicago Press, Chicago*, (1984).
- [16] Birrel, N.D., Davies, P.C.W., Quantum Fields in Curved Spacetime, *The University of Cambridge Press, Cambridge*, (1982).
- [17] Hartman, P., Ordinary Diferantial Equations, *John Wiley & Sons, Inc., New York*, (1964).




Figure 1a. The regions determined by the conditions $\phi_s = \pm \frac{\phi_t}{y}$. Since $\phi_s \rightarrow -\phi_s$ is a discrete symmetry, it is sufficient to work with ϕ in regions *I* and *II*.




Figure 1b. The regions determined by the conditions $\mu_s + \mu = \pm \frac{\mu_t}{y}$. Assuming $1 - 2\mu > 0$, only region *I** is allowed.




Figure 2. General behavior of μ for a parameter value $c = 6\sqrt{\pi}$ in the domain $-q < \phi_s = \psi < p$. The values p , q , $\psi^\pm$ and μ_M are defined by Eqns. (4.10-13).




Figure 3a. Variation of μ with $\psi = \phi_s$ for different values of c , for $0 < \psi < p$.




Figure 3b. Variation of μ with s for different values of c for $-\infty < s < \infty$. For $c \leq 0$, μ is monotone while for $c > 0$ it has a unique maximum μ_M at a finite s . For c large, $\mu(s)$ makes a sharp peak at this critical point, as opposed to the nearly flat variation of μ with respect to ϕ_s as seen in Figure 3a.




Figure 4. The variations of μ , y and r with respect to $\psi = \phi_s$ for $c = 6\sqrt{\pi}$ and $0 < \psi < p$. Recall that for $c \leq 0$, μ is monotone but has a unique maximum for $c > 0$. On the other hand, y is monotone for $c \leq 4\sqrt{\pi}$ but has two critical points for $c > 4\sqrt{\pi}$ while ψ is monotone decreasing for all c . We have chosen the initial value of y as $y_o = 1$. For increasing s , μ rises to its maximum μ_M and decays to zero, while y starts from its initial value reaches its maximum, decays nearly to zero especially for large c and finally rises to ∞ very sharply.




Figure 5. The integral curves of the dynamical system (4.14) for the positive branch in the domain $\mu_s + \mu > 0$ and $\mu < 1/2$. The lines $\mu_s + \mu = 0$ and $\mu_s + 2\mu = 0$ are exact solutions. The half line $\mu_s + \mu = 0$, $\mu < 0$ corresponds to the limit $c \rightarrow -\infty$. For the solution curves in between this half line and $\mu_s + 2\mu = 0$, c ranges from $-\infty$ to zero. For these solutions μ is negative, and monotone increasing and reach the origin as $s \rightarrow \infty$. For c positive, all solution curves reach their maximum value $\mu_M < 1/2$ and arrive at the origin as $s \rightarrow \infty$. Note the sharp changes near the $\mu = 1/2$ line for large c .




Figure 6. The direction fields for the dynamical system (5.1) for the negative branch in the regions D_i , $i = 1, \dots, 5$ for $\mu_s + \mu > 0$ and $\mu < 1/2$.




Figure 7. The integral curves of the dynamical system (5.1) for the negative branch in the domain $\mu_s + \mu > 0$ and $\mu < 1/2$. The line $\mu_s + \mu = 0$ is an exact solution. All solution curves in this region spiral around the point $(1/4, 0)$ as $s \rightarrow \infty$. The curves sketched here are determined by their initial values $(\mu(0), \mu_s(0))$, which are respectively $(-1, 1.001)$, $(-0.6, 0.9)$, $(0.2, 1)$, $(0.4, 1)$, $(0.49, 1)$ respectively from right to left. Note that as one moves to right, the solution curves get very close to the line $\mu = 1/2$ but decay quickly to the level $\mu = 1/4$ making small amplitude oscillations around this level.




Figure 8. The variations of y , μ and ϕ_t/y with respect to s for the dynamical system (5.1) and Eqn.(2.6a). The curves sketched here are determined by their initial value $(y(0), \mu(0), \mu_s(0))$, which is $(1, -0.6, 0.9)$. Note that the solution of ϕ_t/y are obtained from Eqn.(2.6c) for $\phi_s = 0$.